

## Medical Coverage Policy | Pulsed Radiofrequency for the Treatment of Chronic Pain



**EFFECTIVE DATE:** 02|19|2018  
**POLICY LAST UPDATED:** 01|19|2022

### OVERVIEW

Pulsed radiofrequency (RF) facet denervation is used to treat neck and back pain originating in facet joints with degenerative changes. Diagnosis of facet joint pain is confirmed by response to nerve blocks. The goal of facet denervation is long-term pain relief. However, the nerves regenerate and, therefore, repeat procedures may be required.

### MEDICAL CRITERIA

Not applicable

### PRIOR AUTHORIZATION

Not applicable

### POLICY STATEMENT

#### Medicare Advantage Plans

Pulsed radiofrequency denervation for the treatment of chronic spinal/back pain is not covered as the evidence is insufficient to determine the effects of the technology on health outcomes.

#### Commercial Products

Pulsed radiofrequency denervation for the treatment of chronic spinal/back pain is considered not medically necessary as the evidence is insufficient to determine the effects of the technology on health outcomes.

### COVERAGE

Benefits may vary between groups and contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage or Subscriber Agreement for applicable not medically necessary/not covered benefits/coverage.

### BACKGROUND

Pulsed RF consists of short bursts of electrical current of high voltage in the RF range but without heating the tissue enough to cause coagulation. RF is suggested as a possibly safer alternative to thermal RF facet denervation. Temperatures do not exceed 42°C at the probe tip versus temperatures in the 60°C range reached in thermal RF denervation, and tissues may cool between pulses. It is postulated that transmission across small unmyelinated nerve fibers is disrupted but not permanently damaged, while large myelinated fibers are not affected.

Pulsed radiofrequency does not appear to be as effective as conventional radiofrequency denervation, and there is insufficient evidence to evaluate the efficacy of other methods of denervation (eg, alcohol, laser, cryodenervation) for facet joint pain. The evidence is insufficient to determine the effects of the technology on health outcomes. Therefore, these techniques are considered not medically necessary.

### CODING

#### Medicare Advantage Plans and Commercial Products

At this time there are no specific CPT code(s) to describe pulsed radiofrequency. Providers should file using the unlisted code below:

**64999** Unlisted procedure, nervous system

## RELATED POLICIES

Lysis of Epidural Adhesions  
Navigated Transcranial Magnetic Stimulation  
Nerve Graft with Radical Prostatectomy  
Occipital Nerve Stimulation – Insertion  
Percutaneous Electrical Nerve Stimulation and Percutaneous Neuromodulation Therapy  
Peripheral Subcutaneous Field Stimulation  
Sphenopalatine Ganglion Block for Headache

## PUBLISHED

[Provider Update, March 2022](#)  
Provider Update, March 2021  
Provider Update, March 2020  
Provider Update, April 2019  
Provider Update, March 2018  
~~Provider Update, March 2017~~

## REFERENCES

1. Centers for Medicare & Medicaid Services (CMS). Local Coverage Determination (LCD): Facet Joint Injections, Medial Branch Blocks, and Facet Joint Radiofrequency Neurotomy (L35936)
2. Centers for Medicare & Medicaid Services (CMS). Local Coverage Article: Billing and Coding: Facet Joint Injections, Medial Branch Blocks, and Facet Joint Radiofrequency Neurotomy (A57826)
3. Centers for Medicare & Medicaid Services (CMS). Local Coverage Determination (LCD): Pain Management (L33622)
4. Centers for Medicare & Medicaid Services (CMS). Local Coverage Article: Billing and Coding: Pain Management (A52863)
5. Boswell MV, Manchikanti L, Kaye AD, et al. A Best-Evidence Systematic Appraisal of the Diagnostic Accuracy and Utility of Facet (Zygapophysial) Joint Injections in Chronic Spinal Pain. *Pain Physician*. Jul-Aug 2015; 18(4): E497-533. PMID 26218947
6. Falco FJ, Datta S, Manchikanti L, et al. An updated review of the diagnostic utility of cervical facet joint injections. *Pain Physician*. Nov-Dec 2012; 15(6): E807-38. PMID 23159977
7. Falco FJ, Manchikanti L, Datta S, et al. Systematic review of the therapeutic effectiveness of cervical facet joint interventions: an update. *Pain Physician*. Nov-Dec 2012; 15(6): E839-68. PMID 23159978
8. Falco FJ, Manchikanti L, Datta S, et al. An update of the systematic assessment of the diagnostic accuracy of lumbar facet joint nerve blocks. *Pain Physician*. Nov-Dec 2012; 15(6): E869-907. PMID 23159979
9. Falco FJ, Manchikanti L, Datta S, et al. An update of the effectiveness of therapeutic lumbar facet joint interventions. *Pain Physician*. Nov-Dec 2012; 15(6): E909-53. PMID 23159980
10. Cohen SP, Strassels SA, Kurihara C, et al. Randomized study assessing the accuracy of cervical facet joint nerve (medial branch) blocks using different injectate volumes. *Anesthesiology*. Jan 2010; 112(1): 144-52. PMID 19996954
11. Cohen SP, Stojanovic MP, Crooks M, et al. Lumbar zygapophysial (facet) joint radiofrequency denervation success as a function of pain relief during diagnostic medial branch blocks: a multicenter analysis. *Spine J*. May-Jun 2008; 8(3): 498-504. PMID 17662665
12. Pampati S, Cash KA, Manchikanti L. Accuracy of diagnostic lumbar facet joint nerve blocks: a 2-year follow-up of 152 patients diagnosed with controlled diagnostic blocks. *Pain Physician*. Sep-Oct 2009; 12(5): 855-66. PMID 19787011
13. Manchikanti L, Pampati S, Cash KA. Making sense of the accuracy of diagnostic lumbar facet joint nerve blocks: an assessment of the implications of 50% relief, 80% relief, single block, or controlled diagnostic blocks. *Pain Physician*. Mar-Apr 2010; 13(2): 133-43. PMID 20309379
14. Janapala RN, Manchikanti L, Sanapati MR, et al. Efficacy of Radiofrequency Neurotomy in Chronic Low Back Pain: A Systematic Review and Meta-Analysis. *J Pain Res*. 2021; 14: 2859-2891. PMID 34531682

15. Nath S, Nath CA, Pettersson K. Percutaneous lumbar zygapophysial (Facet) joint neurotomy using radiofrequency current, in the management of chronic low back pain: a randomized double-blind trial. *Spine (Phila Pa 1976)*. May 20 2008; 33(12): 1291-7; discussion 1298. PMID 18496338
16. Tekin I, Mirzai H, Ok G, et al. A comparison of conventional and pulsed radiofrequency denervation in the treatment of chronic facet joint pain. *Clin J Pain*. Jul-Aug 2007; 23(6): 524-9. PMID 17575493
17. van Wijk RM, Geurts JW, Wynne HJ, et al. Radiofrequency denervation of lumbar facet joints in the treatment of chronic low back pain: a randomized, double-blind, sham lesion-controlled trial. *Clin J Pain*. Jul-Aug 2005; 21(4):335-44. PMID 15951652
18. Lakemeier S, Lind M, Schultz W, et al. A comparison of intraarticular lumbar facet joint steroid injections and lumbar facet joint radiofrequency denervation in the treatment of low back pain: a randomized, controlled, double-blind trial. *Anesth Analg*. Jul 2013; 117(1): 228-35. PMID 23632051
19. Manchikanti L, Kaye AD, Boswell MV, et al. A Systematic Review and Best Evidence Synthesis of the Effectiveness of Therapeutic Facet Joint Interventions in Managing Chronic Spinal Pain. *Pain Physician*. Jul-Aug 2015; 18(4): E535-82. PMID 26218948
20. Civelek E, Cansever T, Kabatas S, et al. Comparison of effectiveness of facet joint injection and radiofrequency denervation in chronic low back pain. *Turk Neurosurg*. 2012; 22(2): 200-6. PMID 22437295
21. Lord SM, Barnsley L, Wallis BJ, et al. Percutaneous radio-frequency neurotomy for chronic cervical zygapophysial joint pain. *N Engl J Med*. Dec 05 1996; 335(23): 1721-6. PMID 8929263
22. van Eerd M, de Meij N, Kessels A, et al. Efficacy and Long-term Effect of Radiofrequency Denervation in Patients with Clinically Diagnosed Cervical Facet Joint Pain: A Double-blind Randomized Controlled Trial. *Spine (Phila Pa 1976)*. Mar 01 2021; 46(5): 285-293. PMID 33534439
23. Husted DS, Orton D, Schofferman J, et al. Effectiveness of repeated radiofrequency neurotomy for cervical facet joint pain. *J Spinal Disord Tech*. Aug 2008; 21(6): 406-8. PMID 18679094
24. Schofferman J, Kine G. Effectiveness of repeated radiofrequency neurotomy for lumbar facet pain. *Spine (Phila Pa 1976)*. Nov 01 2004; 29(21): 2471-3. PMID 15507813
25. Smuck M, Crisostomo RA, Trivedi K, et al. Success of initial and repeated medial branch neurotomy for zygapophysial joint pain: a systematic review. *PM R*. Sep 2012; 4(9): 686-92. PMID 22980421
26. Rambaransingh B, Stanford G, Burnham R. The effect of repeated zygapophysial joint radiofrequency neurotomy on pain, disability, and improvement duration. *Pain Med*. Sep 2010; 11(9): 1343-7. PMID 20667024
27. Manchikanti L, Singh V, Falco FJ, et al. Comparative outcomes of a 2-year follow-up of cervical medial branch blocks in management of chronic neck pain: a randomized, double-blind controlled trial. *Pain Physician*. Sep-Oct 2010; 13(5): 437-50. PMID 20859313
28. Manchikanti L, Singh V, Falco FJ, et al. Evaluation of lumbar facet joint nerve blocks in managing chronic low back pain: a randomized, double-blind, controlled trial with a 2-year follow-up. *Int J Med Sci*. May 28 2010; 7(3): 124-35. PMID 20567613
29. Manchikanti L, Singh V, Falco FJ, et al. Comparative effectiveness of a one-year follow-up of thoracic medial branch blocks in management of chronic thoracic pain: a randomized, double-blind active controlled trial. *Pain Physician*. Nov-Dec 2010; 13(6): 535-48. PMID 21102966
30. Manchikanti L, Singh V, Falco FJ, et al. The role of thoracic medial branch blocks in managing chronic mid and upper back pain: a randomized, double-blind, active-control trial with a 2-year follow-up. *Anesthesiol Res Pract*. 2012; 2012: 585806. PMID 22851967
31. Hashemi M, Hashemian M, Mohajerani SA, et al. Effect of pulsed radiofrequency in treatment of facet-joint origin back pain in patients with degenerative spondylolisthesis. *Eur Spine J*. Sep 2014; 23(9): 1927-32. PMID 24997616
32. Kroll HR, Kim D, Danic MJ, et al. A randomized, double-blind, prospective study comparing the efficacy of continuous versus pulsed radiofrequency in the treatment of lumbar facet syndrome. *J Clin Anesth*. Nov 2008; 20(7): 534-7. PMID 19041042
33. Van Zundert J, Patijn J, Kessels A, et al. Pulsed radiofrequency adjacent to the cervical dorsal root ganglion in chronic cervical radicular pain: a double blind sham controlled randomized clinical trial. *Pain*. Jan 2007; 127(1-2):173-82. PMID 17055165

34. Iwatsuki K, Yoshimine T, Awazu K. Alternative denervation using laser irradiation in lumbar facet syndrome. *Lasers Surg Med.* Mar 2007; 39(3): 225-9. PMID 17345622
35. Joo YC, Park JY, Kim KH. Comparison of alcohol ablation with repeated thermal radiofrequency ablation in medial branch neurotomy for the treatment of recurrent thoracolumbar facet joint pain. *J Anesth. Jun 2013; 27(3): 390-5.* PMID 23192698
36. Haufe SM, Mork AR. Endoscopic facet debridement for the treatment of facet arthritic pain--a novel new technique. *Int J Med Sci.* May 25 2010; 7(3): 120-3. PMID 20567612
37. Watters WC, Resnick DK, Eck JC, et al. Guideline update for the performance of fusion procedures for degenerative disease of the lumbar spine. Part 13: injection therapies, low-back pain, and lumbar fusion. *J Neurosurg Spine.* Jul 2014; 21(1): 79-90. PMID 24980590
38. Manchikanti L, Kaye AD, Soin A, et al. Comprehensive Evidence-Based Guidelines for Facet Joint Interventions in the Management of Chronic Spinal Pain: American Society of Interventional Pain Physicians (ASIPP) Guidelines Facet Joint Interventions 2020 Guidelines. *Pain Physician.* May 2020; 23(3S): S1-S127. PMID 32503359
39. Cohen SP, Bhaskar A, Bhatia A, et al. Consensus practice guidelines on interventions for lumbar facet joint pain from a multispecialty, international working group. *Reg Anesth Pain Med.* Jun 2020; 45(6): 424-467. PMID 32245841
40. National Institute for Health and Clinical Excellence (NICE). NICE guideline [NG59]: Low back pain and sciatica in over 16s: assessment and management. 2016; <https://www.nice.org.uk/guidance/NG59>. Accessed September 29, 2021.
41. Evidence Based Clinical Guidelines for Multidisciplinary Spine Care: Diagnosis and Treatment of Low Back Pain. American Spine Society. Published 2020. Accessed September 29, 2021.

**CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS**

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

