



EFFECTIVE DATE: 04|01|2017
POLICY LAST UPDATED: 04|19|2023

OVERVIEW

Peripheral subcutaneous field stimulation (PSFS; also called peripheral nerve field stimulation or target field stimulation) is a form of neuromodulation that is intended to treat chronic neuropathic pain.

MEDICAL CRITERIA

Not applicable

PRIOR AUTHORIZATION

Not applicable

POLICY STATEMENT

Medicare Advantage Plans

Peripheral subcutaneous field stimulation, PSFS, is considered not covered as the evidence is insufficient to determine the effects of the technology on health outcomes.

Commercial Products

Peripheral subcutaneous field stimulation, PSFS, is considered not medically necessary as the evidence is insufficient to determine the effects of the technology on health outcomes.

COVERAGE

Benefits may vary between groups and contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage or Subscriber Agreement for applicable not medically necessary/not covered benefits/coverage.

BACKGROUND

Chronic, noncancer pain is responsible for a high burden of illness. Common types of chronic pain are lumbar and cervical back pain, chronic headaches, and abdominal pain. All of these conditions can be challenging to treat. Pharmacologic agents are typically the first-line treatment for chronic pain, and several classes of medications are available. These include analgesics (opioid and nonopioid), antidepressants, anticonvulsants, and muscle relaxants. There also are a variety of nonpharmacologic treatments, including physical therapy, exercise, cognitive-behavioral interventions, acupuncture, chiropractic, and massage.

Neuromodulation is another form of nonpharmacologic therapy that is usually targeted toward patients with chronic pain that is refractory to other modalities. Some forms of neuromodulation, such as transcutaneous electrical nerve stimulation and spinal cord stimulation (SCS), are established methods of chronic pain treatment. Peripheral nerve stimulation, which involves placement of an electrical stimulator on a peripheral nerve, is also used for neuropathic pain originating from peripheral nerves.

PSFS is a modification of peripheral nerve stimulation. In PSFS, leads are placed subcutaneously within the area of maximal pain. The objective of PSFS is to stimulate the region of affected nerves, cutaneous afferents, or the dermatomal distribution of the nerves, which then converge back on the spinal cord. Combined spinal cord stimulation and PSFS is also being evaluated.

Similar to spinal cord stimulation or peripheral nerve stimulation, permanent implantation is preceded by a percutaneous stimulation trial with at least 50% pain reduction. Currently, there is no consensus regarding the indications for PSFS. Criteria for a PSFS trial may include a clearly defined, discrete focal area of pain with a neuropathic or combined somatic/neuropathic pain component with characteristics of burning and increased sensitivity, and failure to respond to other conservative treatments including medications, psychological therapies, physical therapies, surgery, and pain management programs.

The mechanism of action in PSFS is not known. Theories include an increase in endogenous endorphins and other opiate-like substances, modulation of smaller A-delta and C nerve fibers with stimulation of large-diameter A-beta fibers, local stimulation of nerve endings in the skin, local anti-inflammatory and membrane depolarizing effect, or a central action via antegrade activation of A-beta nerve fibers. Complications of PSFS include lead migration or breakage and infection of the lead or neurostimulator.

For individuals who have chronic neuropathic pain who receive PSFS, the evidence includes four ~~two~~ randomized controlled trials (RCTs), a nonrandomized comparative study, and case series. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. One RCT, which used a crossover design, did not compare PSFS with alternatives. Rather, it compared different methods of PSFS. Among trial participants, 24 (80%) of 30 patients had at least a 50% reduction in pain with any type of PSFS. However, because the RCT did not include a sham group or comparator with a different active intervention, this trial offers little evidence for efficacy beyond that of a prospective, uncontrolled study. Another RCT compared sham to external non-invasive peripheral electrical nerve stimulation, but found no significant differences in pain scores between groups after intervention. A third small, pilot RCT found significantly reduced opioid consumption and mean daily pain scores within the first 7 postoperative days in subjects receiving foot, ankle, knee, or shoulder surgery. However, differences in average pain, worst pain, and Defense and Veterans Pain Rating Scale scores were not significantly different between treatment and sham groups following completion of the treatment period on postoperative days 15 and 30. A fourth small, pilot feasibility RCT compared peripheral nerve stimulation with standard medical care to standard medical care alone in veterans undergoing lower extremity amputation. Greater reductions in average phantom limb pain, residual limb pain, and daily opioid consumption were reported through 3 months with the addition of peripheral nerve stimulation. Case series are insufficient to evaluate patient outcomes due to the variable nature of pain and the subjective nature of pain outcome measures. Larger, prospective controlled trials comparing PSFS with placebo or alternative treatment modalities are needed to determine the efficacy of PSFS for chronic pain. The evidence is insufficient to determine the effects of the technology on health outcomes.

CODING

Medicare Advantage Plans and Commercial Products

There is no specific CPT code for this treatment. Report using unlisted CPT code 64999 (Unlisted procedure, nervous system).

RELATED POLICIES:

Percutaneous Electrical Nerve Stimulation and Percutaneous Neuromodulation Therapy
Pulsed Radiofrequency for the Treatment of Chronic Pain
Occipital Nerve Stimulation – Insertion
Lysis of Epidural Adhesions
Sphenopalatine Ganglion Block for Headache
Nerve Graft with Radical Prostatectomy

PUBLISHED

Provider Update, June 2023
Provider Update, September 2022
Provider Update, August 2021
Provider Update, June 2020

REFERENCES

1. McRoberts WP, Wolkowitz R, Meyer DJ, et al. Peripheral nerve field stimulation for the management of localized chronic intractable back pain: results from a randomized controlled study. *Neuromodulation*. Nov-Dec 2013; 16(6): 565-74; discussion 574-5. PMID 23577773
2. Johnson S, Marshall A, Hughes D, et al. Mechanistically informed non-invasive peripheral nerve stimulation for peripheral neuropathic pain: a randomised double-blind sham-controlled trial. *J Transl Med*. Nov 06 2021; 19(1):458. PMID 34742297
3. Mironer YE, Hutcheson JK, Satterthwaite JR, et al. Prospective, two-part study of the interaction between spinal cord stimulation and peripheral nerve field stimulation in patients with low back pain: development of a new spinal-peripheral neurostimulation method. *Neuromodulation*. Mar-Apr 2011; 14(2): 151-4; discussion 155. PMID 21992203
4. Kloimstein H, Likar R, Kern M, et al. Peripheral nerve field stimulation (PNFS) in chronic low back pain: a prospective multicenter study. *Neuromodulation*. Feb 2014; 17(2): 180-7. PMID 24320718
5. Sator-Katzenschlager S, Fiala K, Kress HG, et al. Subcutaneous target stimulation (STS) in chronic noncancer pain: a nationwide retrospective study. *Pain Pract*. Jul-Aug 2010; 10(4): 279-86. PMID 20230450
6. Verrills P, Vivian D, Mitchell B, et al. Peripheral nerve field stimulation for chronic pain: 100 cases and review of the literature. *Pain Med*. Sep 2011; 12(9): 1395-405. PMID 21812906
7. Verrills P, Rose R, Mitchell B, et al. Peripheral nerve field stimulation for chronic headache: 60 cases and long-term follow-up. *Neuromodulation*. Jan 2014; 17(1): 54-9. PMID 24165152
8. Warner NS, Schaefer KK, Eldrige JS, et al. Peripheral Nerve Stimulation and Clinical Outcomes: A Retrospective Case Series. *Pain Pract*. Apr 2021; 21(4): 411-418. PMID 33222402
9. National Institute for Health and Care Excellence (NICE). Peripheral nerve-field stimulation for chronic low back pain [IPG451]. 2013; <https://www.nice.org.uk/guidance/ipg451>. Accessed March 28, 2023.
10. Albright-Trainer B, Phan T, Trainer RJ, et al. Peripheral nerve stimulation for the management of acute and subacute post-amputation pain: a randomized, controlled feasibility trial. *Pain Manag*. Apr 2022; 12(3): 357-369. PMID 34761694
11. Strand N, D'Souza RS, Hagedorn JM, et al. Evidence-Based Clinical Guidelines from the American Society of Pain and Neuroscience for the Use of Implantable Peripheral Nerve Stimulation in the Treatment of Chronic Pain. *J Pain Res*. 2022; 15: 2483-2504. PMID 36039168
12. Ilfeld BM, Plunkett A, Vijeswarapu AM, et al. Percutaneous Peripheral Nerve Stimulation (Neuromodulation) for Postoperative Pain: A Randomized, Sham-controlled Pilot Study. *Anesthesiology*. Jul 01 2021; 135(1): 95-110. PMID 33856424

CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

