

**Medical Coverage Policy | Bone Turnover Markers for the Diagnosis and Management of Osteoporosis and Diseases Associated With High Bone Turnover**



**EFFECTIVE DATE:** 10|1|2015

**POLICY LAST UPDATED:** 7|7|2015

### **OVERVIEW**

Bone turnover markers are biochemical markers of either bone formation or bone resorption. Commercially available tests are available to assess some of these markers in urine and/or serum by high performance liquid chromatography or immunoassay. Assessment of bone turnover markers is proposed to supplement bone mineral density (BMD) measurement in the diagnosis of osteoporosis and to aid in treatment decisions. Bone turnover markers could also potentially be used to evaluate treatment effectiveness before changes in BMD can be observed.

### **MEDICAL CRITERIA**

Not applicable

### **PRIOR AUTHORIZATION**

Prior authorization review is not required.

### **POLICY STATEMENT**

#### **BlueCHiP for Medicare and Commercial Products**

Measurement of bone turnover markers is considered not medically necessary in the diagnosis and management of osteoporosis or in the management of patients with conditions associated with high rates of bone turnover, including but not limited to Paget disease, primary hyperparathyroidism, and renal osteodystrophy.

### **COVERAGE**

Benefits may vary between groups and contracts. Please refer to the appropriate Evidence of Coverage or Subscriber Agreement for applicable not medically necessary benefits/coverage.

### **BACKGROUND**

After cessation of growth, bone is in a constant state of remodeling (or turnover), with initial absorption of bone by osteoclasts followed by deposition of new bone matrix by osteoblasts. This constant bone turnover is critical to the overall health of the bone, by repairing microfractures and remodeling the bony architecture in response to stress. Normally, the action of osteoclasts and osteoblasts is balanced, but bone loss occurs if the two processes become uncoupled. Bone turnover markers can be categorized as bone formation markers or bone resorption markers and can be identified in serum and/or urine.

There is interest in the use of bone turnover markers to evaluate age-related osteoporosis, a condition characterized by slow, prolonged bone loss, resulting in an increased risk of fractures at the hip, spine, or wrist. Currently, fracture risk is primarily based on measurements of BMD in conjunction with other genetic and environmental factors, such as family history of osteoporosis, history of smoking, and weight. It is thought that the level of bone turnover markers may also predict fracture risk, possibly through a different mechanism than that associated with BMD. However, it must be emphasized that the presence of bone turnover markers in the serum or urine is not necessarily related to bone loss. For example, even if bone turnover is high, if resorption is balanced with formation, there will be no net bone loss. Bone loss will only occur if resorption exceeds formation. Therefore, bone turnover markers have been primarily studied as an adjunct, not an alternative, to measurements of BMD to estimate fracture risk and document the need for preventive or therapeutic strategies for osteoporosis.

In addition, bone turnover markers might provide a more immediate assessment of treatment response and predict change in BMD in response to treatment. Treatment-related changes in BMD occur very slowly. This fact, coupled with the precision of BMD technologies, suggested that clinically significant changes in BMD could not be reliably detected until at least 2 years. In contrast, changes in bone turnover markers could be anticipated after 3 months of therapy. Bone turnover markers have also been researched as markers of diseases associated with markedly high levels of bone turnover, such as Paget disease, primary hyperparathyroidism, and renal osteodystrophy.

The literature suggests that bone turnover marker levels may be independently associated with osteoporosis and fracture risk in some groups, but there is insufficient evidence reporting an association for any specific marker. Questions remain about whether bone turnover markers are sufficiently sensitive to reliably determine individual treatment responses. In addition, there is insufficient evidence from controlled studies that bone turnover marker measurement improves adherence to treatment, impacts management decisions, and/or improves health outcomes such as reducing fracture rates. Thus, the use of bone turnover markers for the diagnosis and management of osteoporosis is considered not medically necessary.

There is insufficient evidence that measurement of bone turnover markers improves patient management or health outcomes in patients with conditions associated with high bone turnover including Paget disease, primary hyperparathyroidism, and renal osteodystrophy. Thus, bone turnover marker testing for these other conditions is considered not medically necessary.

#### **CODING**

The following CPT codes are medically necessary when filed with a covered diagnosis:

**82523** Collagen cross links, any method

**83937** Osteocalcin (bone gla protein)

The following is a list of diagnosis codes that are not medically necessary for this service:



Bone Turnover  
Markers for Diagnosis

#### **RELATED POLICIES**

None

#### **PUBLISHED**

Provider Update, August 2015

#### **REFERENCES**

1. Tamaki J, Iki M, Kadowaki E, et al. Biochemical markers for bone turnover predict risk of vertebral fractures in postmenopausal women over 10 years: the Japanese Population-based Osteoporosis (JPOS) Cohort Study. *Osteoporos Int.* Mar 2013;24(3):887-897. PMID 22885773
2. Bauer DC, Garnero P, Harrison SL, et al. Biochemical markers of bone turnover, hip bone loss and fracture in older men: the MrOS Study. *J Bone Mineral Res.* 2009;24(12):2032-2038. PMID
3. Johansson H, Oden A, Kanis JA, et al. A meta-analysis of reference markers of bone turnover for prediction of fracture. *Calcif Tissue Int.* May 2014;94(5):560-567. PMID 24590144
4. Biver E, Chopin F, Coiffier G, et al. Bone turnover markers for osteoporotic status assessment? A systematic review of their diagnosis value at baseline in osteoporosis. *Joint Bone Spine.* Jan 2012;79(1):20-25. PMID 21724445

5. Bauer DC, Garnero P, Hochberg MC, et al. Pretreatment levels of bone turnover and the antifracture efficacy of alendronate: the Fracture Intervention trial. *J Bone Miner Res.* 2006;21(2):292-299. PMID
6. Abe Y, Ishikawa H, Fukao A. Higher efficacy of urinary bone resorption marker measurements in assessing response to treatment for osteoporosis in postmenopausal women. *Tohoku J Exp Med.* 2008;214(1):51-59. PMID
7. Shiraki M, Itabashi A. Short-term menatetrenone therapy increases gamma-carboxylation of osteocalcin with a moderate increase of bone turnover in postmenopausal osteoporosis: a randomized prospective study. *J Bone Miner Metab.* 2009;27(3):333-340. PMID
8. Funck-Brentano T, Biver E, Chopin F, et al. Clinical utility of serum bone turnover markers in postmenopausal osteoporosis therapy monitoring: a systematic review. *Semin Arthritis Rheum.* Oct 2011;41(2):157-169. PMID 21507464
9. Baxter I, Rogers A, Eastell R, et al. Evaluation of urinary N-telopeptide of type I collagen measurements in the management of osteoporosis in clinical practice. *Osteoporos Int.* Mar 2013;24(3):941-947. PMID 22872068
10. National Osteoporosis Foundation. 2014 Clinician's guide to prevention and treatment of osteoporosis. <http://nof.org/files/nof/public/content/file/2610/upload/895.pdf>. Accessed July, 2014
11. Burch J, Rice S, Yang H, et al. Systematic review of the use of bone turnover markers for monitoring the response to osteoporosis treatment: the secondary prevention of fractures, and primary prevention of fractures in high-risk groups. *Health Technol Assess.* Feb 2014;18(11):1-180. PMID 24534414
12. Roux C, Giraudeau B, Rouanet S, et al. Monitoring of bone turnover markers does not improve persistence with ibandronate treatment. *Joint Bone Spine* 2011;79(4):389-392. PMID
13. Rianon N, Alex G, Callender G, et al. Preoperative serum osteocalcin may predict postoperative elevated parathyroid hormone in patients with primary hyperparathyroidism. *World J Surg.* Jun 2012;36(6):1320-1326. PMID 22278606
14. Woitge HW, Oberwittler H, Heichel S, et al. Short- and long-term effects of ibandronate treatment on bone turnover in Paget disease of bone. *Clin Chem.* May 2000;46(5):684-690. PMID 10794751
15. Reid IR, Davidson JS, Wattie D, et al. Comparative responses of bone turnover markers to bisphosphonate therapy in Paget's disease of bone. *Bone.* Jul 2004;35(1):224-230. PMID 15207761
16. Alvarez L, Guanabens N, Peris P, et al. Usefulness of biochemical markers of bone turnover in assessing response to the treatment of Paget's disease. *Bone.* Nov 2001;29(5):447-452. PMID 11704497
17. North American Menopause Society. Management of osteoporosis in postmenopausal women: 2010 position statement of the North American Menopause Society.
18. Vasikaran S, Cooper C, Eastell R, et al. International Osteoporosis Foundation and International Federation of Clinical Chemistry and Laboratory Medicine Position on bone marker standards in osteoporosis. *Clin Chem Lab Med.* 2011;49(8):1271-1274. PMID
19. McCloskey EV, Vasikaran S, Cooper C. Official Positions for FRAX(R) clinical regarding biochemical markers from Joint Official Positions Development Conference of the International Society for Clinical Densitometry and International Osteoporosis Foundation on FRAX(R). *J Clin Densitom.* Jul-Sep 2011;14(3):220-222. PMID 21810528

20. U.S. Preventive Services Task Force (USPSTF). Screening for Osteoporosis. 2011; <http://www.uspreventiveservicestaskforce.org/uspstf10/osteoporosis/osteosum.htm>. Accessed July, 2014.
21. CMS National Coverage Determinations. 190.19 Collagen Crosslinks, Any Method. <http://www.cms.hhs.gov/CoverageGenInfo/Downloads/manual200907.pdf>. Accessed July, 2014.
22. Federal Register, November 23, 2001. Rules and Regulations.66(No. 226). PMID

**CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS**

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

