

**EFFECTIVE DATE:** ||  
**POLICY LAST UPDATED:** 06|23|2015

## OVERVIEW

Image-guided minimally invasive lumbar decompression (IG-MLD) describes a novel percutaneous procedure for decompression of the central spinal canal in patients with lumbar spinal stenosis (LSS). In this procedure, a specialized cannula and surgical tools (mild®) are used under fluoroscopic guidance for bone and tissue sculpting near the spinal canal.

## MEDICAL CRITERIA

Not applicable

## PRIOR AUTHORIZATION

Not applicable

## POLICY STATEMENT

### BlueCHiP for Medicare

Percutaneous image-guided lumbar decompression for lumbar spinal stenosis (PILD) is covered only for members enrolled in a Medicare approved clinical trial under section 1862(a)(1)(E) through Coverage with Evidence Development (CED) for beneficiaries with LSS who are enrolled in an approved clinical study that meets CMS (Centers for Medicare and Medicaid Services) criteria. Clinical trials may be found at <http://www.clinicaltrials.gov/>

Medicare policy is developed separately from BCBSRI policy. Medicare policy incorporates consideration of governmental regulations from CMS, such as national coverage determinations or local coverage determinations. In addition to benefit differences, CMS may reach different conclusions regarding the scientific evidence than does BCBSRI. Medicare and BCBSRI policies may differ. However, BlueCHiP for Medicare members must be offered, at least, the same services as Medicare offers.

### Commercial products

Image-guided minimally invasive lumbar decompression is considered not medically necessary as there is insufficient peer reviewed literature that demonstrates that the procedure is effective.

## COVERAGE

Benefits may vary between groups/contracts. Please refer to the appropriate Evidence of Coverage or Subscriber Agreement for limitations of benefits/coverage when services are not medically necessary.

## BACKGROUND

In LSS, the space around the spinal cord narrows, compressing the spinal cord and the nerve roots. The most common symptom of LSS is back pain with neurogenic claudication, i.e., pain, numbness, or weakness in the legs that worsens with standing or walking and is alleviated with sitting or leaning forward. Compression of neural elements generally occurs from a combination of degenerative changes including ligamentum flavum hypertrophy, bulging of the intervertebral disc, and facet thickening with arthropathy. Spinal stenosis is often linked to age-related changes in disc height and arthritis of the facet joints. LSS is one of the most common reasons for back surgery and the most common reason for lumbar spine surgery in adults older than 65 years of age. The goal of surgical treatment is to “decompress” the spinal cord and/or nerve roots. For patients with LSS, surgical laminectomy has established benefits in reducing pain and improving quality of life. Less

invasive surgical procedures have been developed, such as open laminotomy and microendoscopic laminotomy. Limited evidence on the comparative efficacy of these procedures suggests that less invasive procedures may achieve a roughly similar benefit with less adverse effects. The present policy addresses posterior decompression of central LSS with a percutaneous treatment that is performed under fluoroscopic guidance. Percutaneous IG-MLD using a specially designed tool kit (mild®) has been proposed as an ultraminimally invasive treatment of central LSS. In this procedure, the epidural space is filled with contrast medium under fluoroscopic guidance. Using a 6-gauge cannula that is clamped in place with a back plate, single-use tools (portal cannula, surgical guide, bone rongeur, tissue sculpter, trocar) are used to resect thickened ligamentum flavum and small pieces of lamina. The tissue and bone sculpting is conducted entirely under fluoroscopic guidance, with additional contrast media added throughout the procedure to aid visualization of the decompression. The process is repeated on the opposite side for bilateral decompression of the central canal. The devices are not intended to be used near the lateral neural elements and are contraindicated for disc procedures.

Alternative posterior decompressive surgical procedures include:

- Decompressive laminectomy, the classic treatment for LSS, which unroofs the spinal canal by extensive resection of posterior spinal elements, including the lamina, spinous processes, portions of the facet joints, ligamentum flavum, and the interspinous ligaments. Wide muscular dissection and retraction is needed to achieve adequate surgical visualization. The extensive resection and injury to the posterior spine and supporting muscles can lead to instability with significant morbidity, both postoperatively and longer term. Spinal fusion, performed at the same time as laminectomy or after symptoms have developed, may be required to reduce the resultant instability. Laminectomy may be used for extensive multilevel decompression.
- Hemilaminotomy and laminotomy, sometimes termed laminoforaminotomy, are less invasive than laminectomy. These procedures focus on the interlaminar space, where most of the pathologic changes are concentrated, minimizing resection of the stabilizing posterior spine. A laminotomy typically removes the inferior aspect of the cranial lamina, superior aspect of the subjacent lamina, ligamentum flavum, and the medial aspect of the facet joint. In contrast to laminectomy, laminotomy does not disrupt the facet joints, supra- and interspinous ligaments, a major portion of the lamina, or the muscular attachments. Muscular dissection and retraction are required to achieve adequate surgical visualization.
- Microendoscopic decompressive laminotomy (MEDL) is similar to laminotomy but uses endoscopic visualization. The position of the tubular working channel is confirmed by fluoroscopic guidance, and serial dilators (METRx™ lumbar endoscopic system; Medtronic) are used to dilate the musculature and expand the fascia. For MEDL, an endoscopic curette, rongeur, and drill are used for the laminotomy, facetectomy, and foraminotomy. The working channel may be repositioned from a single incision for multilevel and bilateral dissections.

#### Regulatory Status

The mild® tool kit (Vertos Medical) initially received 510(k) marketing clearance as the X-Sten MILD Tool Kit (X-Sten Corp.) from FDA in 2006, with intended use as a set of specialized surgical instruments to be used to perform percutaneous lumbar decompressive procedures for the treatment of various spinal conditions. Vertos' mild® instructions for use state that the devices are not intended for disc procedures but rather for tissue resection at the perilaminar space, within the interlaminar space, and at the ventral aspect of the lamina. These devices are not intended for use near the lateral neural elements and remain dorsal to the dura using image guidance and anatomical landmarks. Note: The abbreviation MILD has also been used for microscopic image-preserving interlaminar decompression, which involves a small skin incision at the interspinous level and partial drilling of the spinous process, with decompression performed under microscopic visualization.

Posterior decompression for lumbar spinal stenosis (LSS) has been evolving toward increasingly minimally invasive procedures in an attempt to minimize postoperative morbidity and spinal instability. In general, the literature comparing surgical procedures is limited. The evidence available suggests that less invasive surgical decompression may reduce perioperative morbidity without impairing long-term outcomes when performed in appropriately selected patients.

In contrast to conventional surgical decompression, the mild® procedure is a percutaneous decompressive procedure performed solely under fluoroscopic guidance (eg, without endoscopic or microscopic visualization of the work area). This procedure is indicated for central stenosis only, without the capability of addressing nerve root compression or disc herniation, should it be required. One small controlled trial with short-term follow-up and small case series of patients treated with image-guided minimally invasive lumbar decompression report improvements in pain and functioning, but controlled trials are lacking, and the efficacy of this procedure compared with alternatives cannot be determined at this time. Due to the unknown impact on health outcomes, randomized controlled trials in appropriate patients are needed to compare this novel procedure with the established alternatives. Therefore, this procedure is considered not medically necessary as there is insufficient peer-reviewed literature that demonstrates that the procedure is effective.

#### **CODING**

##### **BlueCHiP for Medicare and Commercial products**

The following codes are covered for BlueCHiP for Medicare when filed with the Q0 modifier. These codes are considered not medically necessary for Commercial products.

**0275T**

**G0276**

**Modifier Q0** Investigational clinical service provided in a clinical research study that is in an approved research study

**Note:** Medicare claims filed without the Q0 modifier will deny as not medically necessary.

#### **RELATED POLICIES**

CPT Category III Codes

#### **PUBLISHED**

Provider Update, 2015

#### **REFERENCES:**

1. Chou R, Baisden J, Carragee EJ, et al. Surgery for low back pain: a review of the evidence for an American Pain Society Clinical Practice Guideline. *Spine*. May 1 2009;34(10):1094-1109. PMID 19363455
2. Chou R, Loeser JD, Owens DK, et al. Interventional therapies, surgery, and interdisciplinary rehabilitation for low back pain: an evidence-based clinical practice guideline from the American Pain Society. *Spine*. May 1 2009;34(10):1066-1077. PMID 19363457
3. Weinstein JN, Lurie JD, Tosteson TD, et al. Surgical versus nonsurgical treatment for lumbar degenerative spondylolisthesis. *N Engl J Med*. May 31 2007;356(22):2257-2270. PMID 17538085
4. Weinstein JN, Tosteson TD, Lurie JD, et al. Surgical versus nonsurgical therapy for lumbar spinal stenosis. *N Engl J Med*. Feb 21 2008;358(8):794-810. PMID 18287602
5. Kreiner DS, Macvicar J, Duszynski B, et al. The mild(R) Procedure: A Systematic Review of the Current Literature. *Pain Med*. Feb 2014;15(2):196-205. PMID 24308292

6. Chopko BW. Long-term results of percutaneous lumbar decompression for LSS: two-year outcomes. Clin J Pain. Nov 2013;29(11):939-943. PMID 23446067
7. Brown LL. A double-blind, randomized, prospective study of epidural steroid injection vs. the mild(R) procedure in patients with symptomatic lumbar spinal stenosis. Pain Pract. Jun 2012;12(5):333-341. PMID 22272730
8. Chopko B, Caraway DL. MiDAS I (mild Decompression Alternative to Open Surgery): a preliminary report of a prospective, multi-center clinical study. Pain Physician. Jul-Aug 2010;13(4):369-378. PMID 20648206
9. Mekhail N, Vallejo R, Coleman MH, et al. Long-term results of percutaneous lumbar decompression mild((R)) for spinal stenosis. Pain Pract. Mar 2012;12(3):184-193. PMID 21676166

[CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS](#)

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

