

Medical Coverage Policy | Ultrasonographic Measurement of Carotid Intima-Medial Thickness as an Assessment of subclinical Atherosclerosis Dynamic Posturagraphy



EFFECTIVE DATE: 10|1|2015
POLICY LAST UPDATED: 7|7|2016

OVERVIEW

Ultrasonographic measurement of carotid intima-medial (or intimal-media) thickness (CIMT) refers to the use of B mode ultrasound to determine the thickness of the 2 innermost layers of the carotid artery wall, the intima and the media. Detection and monitoring of intima-medial thickening, which is a surrogate marker for atherosclerosis, may provide an opportunity to intervene earlier in atherogenic disease and/or monitor disease progression.

MEDICAL CRITERIA

Not applicable

PRIOR AUTHORIZATION

Not applicable

POLICY STATEMENT

BlueCHiP for Medicare and Commercial Products

Ultrasonographic measurement of carotid artery intima-medial thickness (CIMT) as a technique of identifying subclinical atherosclerosis is considered not medically necessary for use in the screening, diagnosis, or management of atherosclerotic disease. The existing data are insufficient to determine the impact of this technology on net health outcome.

COVERAGE

Benefits may vary between groups and contracts. Please refer to the appropriate Evidence of Coverage or Subscriber Agreement for applicable not medically necessary benefits/coverage.

BACKGROUND

Coronary heart disease (CHD) accounts for 27% of all deaths in the U.S.(1) Established major risk factors for CHD have been identified by the National Cholesterol Education Program Expert Panel (NCEP). These risk factors include elevated serum levels of low density lipoprotein (LDL) cholesterol, total cholesterol, and reduced levels of high density lipoprotein (HDL) cholesterol. Other risk factors include a history of cigarette smoking, hypertension, family history of premature CHD, and age.

The third report of the NCEP Adult Treatment Panel (ATP III) establishes various treatment strategies to modify the risk of CHD, with emphasis on target goals of LDL cholesterol. Pathology studies have demonstrated that levels of traditional risk factors are associated with the extent and severity of atherosclerosis. ATP III recommends use of the Framingham criteria to further stratify those patients with 2 or more risk factors for more intensive lipid management.(2) However, at every level of risk factor exposure, there is substantial variation in the amount of atherosclerosis, presumably related to genetic susceptibility and the influence of other risk factors. Therefore, there has been interest in identifying a technique that can improve the ability to diagnose those at risk of developing CHD, as well as measure disease progression, particularly for those at intermediate risk.

The carotid arteries can be well-visualized by ultrasonography, and ultrasonographic measurement of the CIMT has been investigated as a technique to identify and monitor subclinical atherosclerosis. B mode ultrasound is most commonly used to measure CIMT. The intima-medial thickness (IMT) is measured and averaged over several sites in each carotid artery. Imaging of the far wall of each common carotid artery yields more accurate and reproducible IMT measurements than imaging of the near wall. Two echogenic lines are produced, representing the lumen-intima interface and the media-adventitia interface. The distance between these 2 lines constitutes the IMT.

In addition, available studies do not define how the use of CIMT in clinical practice improves outcomes. There appears to be no scientific literature that directly and experimentally tests the hypothesis that measurement of CIMT results in improved patient outcomes and no specific guidance on how measurements of CIMT should be incorporated into risk assessment and risk management. The existing data are insufficient to determine the impact of this technology on net health outcome. Therefore, CIMT is considered investigational for use in the screening, diagnosis, or management of atherosclerotic disease.

Regulatory Status

In February 2003, SonoCalc® (SonoMetric Health, LLC, Bountiful, UT) was cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process. The FDA determined that this software was substantially equivalent to existing image display products for use in the automatic measurement of the IMT of the carotid artery from images obtained from ultrasound systems. Subsequently, several other devices have been approved through the 510(k) process. Product code: LLZ.

CODING

The following codes are not medically necessary:

93895 Quantitative carotid intima media thickness and carotid atheroma evaluation, bilateral (new code 01/01/15)

0126T Common carotid intima-medial thickness (IMT) study for evaluation of atherosclerotic burden or coronary heart disease risk factor assessment

RELATED POLICIES

None

PUBLISHED

Provider Update, August 2015

REFERENCES

1. Minino AM, Heron MP, Murphy SL et al. Deaths: final data for 2004. Natl Vital Stat Rep 2007; 55(19):1-119.
2. Pasternak RC. Report of the Adult Treatment Panel III: the 2001 National Cholesterol Education Program guidelines on the detection, evaluation and treatment of elevated cholesterol in adults. *Cardiol Clin* 2003; 21(3):393-8.
3. Den Ruijter HM, Peters SA, Anderson TJ et al. Common carotid intima-media thickness measurements in cardiovascular risk prediction: a meta-analysis. *JAMA* 2012; 308(8):796-803.
4. van den Oord SC, Sijbrands EJ, ten Kate GL et al. Carotid intima-media thickness for cardiovascular risk assessment: systematic review and meta-analysis. *Atherosclerosis* 2013; 228(1):1-11.
5. Lorenz MW, Polak JF, Kavousi M et al. Carotid intima-media thickness progression to predict cardiovascular events in the general population (the PROG-IMT collaborative project): a metaanalysis of individual participant data. *Lancet* 2012; 379(9831):2053-62.

6. Peters SA, den Ruijter HM, Bots ML et al. Improvements in risk stratification for the occurrence of cardiovascular disease by imaging subclinical atherosclerosis: a systematic review. *Heart* 2012; 98(3):177-84.
7. Plichart M, Celermajer DS, Zureik M et al. Carotid intima-media thickness in plaque-free site, carotid plaques and coronary heart disease risk prediction in older adults. The Three-City Study. *Atherosclerosis* 2011; 219(2):917-24.
8. Keo HH, Baumgartner I, Hirsch AT et al. Carotid plaque and intima-media thickness and the incidence of ischemic events in patients with atherosclerotic vascular disease. *Vasc Med* 2011; 16(5):323-30.
9. Nambi V, Chambless L, He M et al. Common carotid artery intima-media thickness is as good as carotid intima-media thickness of all carotid artery segments in improving prediction of coronary heart disease risk in the Atherosclerosis Risk in Communities (ARIC) study. *Eur Heart J* 2012; 33(2):183-90.
10. Xie W, Liang L, Zhao L et al. Combination of carotid intima-media thickness and plaque for better predicting risk of ischaemic cardiovascular events. *Heart* 2011; 97(16):1326-31.
11. Mookadam F, Moustafa SE, Lester SJ et al. Subclinical atherosclerosis: evolving role of carotid intima-media thickness. *Prev Cardiol* 2010; 13(4):186-97.
12. Dobs AS, Nieto FJ, Szklo M et al. Risk factors for popliteal and carotid wall thicknesses in the Atherosclerosis Risk in Communities (ARIC) Study. *Am J Epidemiol* 1999; 150(10):1055-67.
13. Chambless LE, Heiss G, Folsom AR et al. Association of coronary heart disease incidence with carotid arterial wall thickness and major risk factors: the Atherosclerosis Risk in Communities (ARIC) Study, 1987-1993. *Am J Epidemiol* 1997; 146(6):483-94.
14. van der Meer IM, Bots ML, Hofman A et al. Predictive value of noninvasive measures of atherosclerosis for incident myocardial infarction: the Rotterdam Study. *Circulation* 2004; 109(9):1089-94.
15. O'Leary DH, Polak JF, Kronmal RA et al. Carotid-artery intima and media thickness as a risk factor for myocardial infarction and stroke in older adults. Cardiovascular Health Study Collaborative Research Group. *N Engl J Med* 1999; 340(1):14-22.
16. Lorenz MW, Schaefer C, Steinmetz H et al. Is carotid intima media thickness useful for individual prediction of cardiovascular risk? Ten-year results from the Carotid Atherosclerosis Progression Study (CAPS). *Eur Heart J* 2010; 31(16):2041-8.
17. Probstfield JL, Margitic SE, Byington RP et al. Results of the primary outcome measure and clinical events from the Asymptomatic Carotid Artery Progression Study. *Am J Cardiol* 1995; 76(9):47C-53C.
18. Byington RP, Evans GW, Espeland MA et al. Effects of lovastatin and warfarin on early carotid atherosclerosis: sex-specific analyses. Asymptomatic Carotid Artery Progression Study (ACAPS) Research Group. *Circulation* 1999; 100(3):e14-7.
19. Hodis HN, Mack WJ, LaBree L et al. Reduction in carotid arterial wall thickness using lovastatin and dietary therapy: a randomized controlled clinical trial. *Ann Intern Med* 1996; 124(6):548-56.
20. Folsom AR, Kronmal RA, Detrano RC et al. Coronary artery calcification compared with carotid intima-media thickness in the prediction of cardiovascular disease incidence: the Multi-Ethnic Study of Atherosclerosis (MESA). *Arch Intern Med* 2008; 168(12):1333-9.

21. Paramsothy P, Knopp RH, Bertoni AG et al. Association of combinations of lipid parameters with carotid intima-media thickness and coronary artery calcium in the MESA (Multi-Ethnic Study of Atherosclerosis). J Am Coll Cardiol 2010; 56(13):1034-41.

22. Blaha MJ, Rivera JJ, Budoff MJ et al. Association between obesity, high-sensitivity C-reactive protein ≥ 2 mg/L, and subclinical atherosclerosis: implications of JUPITER from the Multi-Ethnic Study of Atherosclerosis. Arterioscler Thromb Vasc Biol 2011; 31(6):1430-8.

[CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS](#)

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

