

Medical Coverage Policy



**Blue Cross
Blue Shield**
of Rhode Island

Manipulation Under Anesthesia

Device/Equipment Drug Medical Surgery Test Other

Effective Date:	5/18/2010	Policy Last Updated:	3/19/2013
------------------------	------------------	-----------------------------	------------------

Prospective review is recommended/required. Please check the member agreement for preauthorization guidelines.

Prospective review is not required.

Description:

Manipulation under anesthesia (MUA) consists of a series of mobilization, stretching, and traction procedures performed while the patient receives anesthesia (usually general anesthesia or moderate sedation). This policy does not address manipulation under anesthesia for fractures, completely dislocated joints, adhesive capsulitis (e.g., frozen shoulder), and/or fibrosis of a joint that may occur following total joint replacement.

Manipulation is intended to break up fibrous and scar tissue to relieve pain and improve range of motion. Anesthesia or sedation is used to reduce pain, spasm, and reflex muscle guarding that may interfere with the delivery of therapies and to allow the therapist to break up joint and soft-tissue adhesions with less force than would be required to overcome patient resistance or apprehension. MUA is generally performed with an anesthesiologist in attendance. Manipulation has also been performed after injection of local anesthetic into lumbar zygapophyseal and/or sacroiliac joints under fluoroscopic guidance (MUJA) and after epidural injection of corticosteroid and local anesthetic (MUESI).

MUA is described as follows: after sedation is achieved, a series of mobilization, stretching, and traction procedures to the spine and lower extremities is performed and may include passive stretching of the gluteal and hamstring muscles with straight leg raise, hip capsule stretching and mobilization, lumbosacral traction, and stretching of the lateral abdominal and paraspinal muscles. After the stretching and traction procedures, spinal manipulative therapy (SMT) is delivered with high-velocity, short-amplitude thrust applied to a spinous process by hand while the upper torso and lower extremities are stabilized. SMT may also be applied to the thoracolumbar or cervical area if considered necessary to address the low back pain. The MUA takes 15–20 minutes, and after recovery from anesthesia the patient is discharged with instructions to remain active and use heat or ice for short-term analgesic control. Some practitioners recommend performing the procedure on 3 consecutive days for best results. Care after MUA may include 4–8 weeks of active rehabilitation with manual therapy including SMT and other modalities.

Scientific evidence regarding spinal manipulation under anesthesia, spinal manipulation with joint anesthesia, and spinal manipulation after epidural anesthesia and corticosteroid injection is limited to observational case series and nonrandomized comparative studies. These data are insufficient to determine whether MUA improves health outcomes, thus it is considered not medically necessary.

Medical Criteria:

None

Policy

Spinal manipulation (and manipulation of other joints, e.g., hip joint, performed during the procedure) with the patient under anesthesia, spinal manipulation under joint anesthesia, and spinal manipulation after epidural anesthesia and corticosteroid injection are considered not medically necessary for treatment of chronic spinal (cranial, cervical, thoracic, lumbar) pain and chronic sacroiliac and pelvic pain as there is insufficient peer-reviewed scientific literature that demonstrates that the procedure/service is effective.

Coverage:

Benefits may vary by group/contract. Please refer to the appropriate member certificate/subscriber agreement for services not medically necessary benefits/coverage.

Coding

The following codes are not medically necessary when used for the indications above.

22505, 00640

Related Topics:

None

Published:

Provider Update, June 2013

Provider Update, May 2012

Provider Update, July 2011

Provider Update, July 2010

References:

- I. Dagenais S, Mayer J, Wooley JR et al. Evidence-informed management of chronic low back pain with medicine-assisted manipulation. *Spine J* 2008; 8(1):142-91
- II. West DT, Mathews RS, Miller MR et al. Effective management of spinal pain in one hundred seventy-seven patients evaluated for manipulation under anesthesia. *J Manipulative Physiol Ther* 1999; 22(5):299-308.
- III. Kohlbeck FJ, Haldeman S, Hurwitz EL et al. Supplemental care with medication-assisted manipulation versus spinal manipulation therapy alone for patients with chronic low back pain. *J Manipulative Physiol Ther* 2005; 28(4):245-52.

History:

February 2013 - Annual review

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are

determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice.